



Ecdysozoan phylogeny and Bayesian inference: first use of nearly complete 28S and 18S rRNA gene sequences to classify the arthropods and their kin[☆]

Jon M. Mallatt,^{a,*} James R. Garey,^b and Jeffrey W. Shultz^c

^a School of Biological Sciences, Washington State University, Pullman, WA 99164-4236, USA

^b Department of Biology, University of South Florida, 4202 East Fowler Ave. SCA110, Tampa, FL 33620, USA

^c Department of Entomology, University of Maryland, College Park, MD 20742, USA

Received 4 March 2003; revised 18 July 2003

Abstract

Relationships among the ecdysozoans, or molting animals, have been difficult to resolve. Here, we use nearly complete 28S + 18S ribosomal RNA gene sequences to estimate the relations of 35 ecdysozoan taxa, including newly obtained 28S sequences from 25 of these. The tree-building algorithms were likelihood-based Bayesian inference and minimum-evolution analysis of LogDet-transformed distances, and hypotheses were tested with parametric bootstrapping. Better taxonomic resolution and recovery of established taxa were obtained here, especially with Bayesian inference, than in previous parsimony-based studies that used 18S rRNA sequences (or 18S plus small parts of 28S). In our gene trees, priapulid worms represent the basal ecdysozoans, followed by nematomorphs, or nematomorphs plus nematodes, followed by Panarthropoda. Panarthropoda was monophyletic with high support, although the relationships among its three phyla (arthropods, onychophorans, tardigrades) remain uncertain. The four groups of arthropods—hexapods (insects and related forms), crustaceans, chelicerates (spiders, scorpions, horseshoe crabs), and myriapods (centipedes, millipedes, and relatives)—formed two well-supported clades: Hexapoda in a paraphyletic crustacea (Pancrustacea), and ‘Chelicerata + Myriapoda’ (a clade that we name ‘Paradoxopoda’). Pycnogonids (sea spiders) were either chelicerates or part of the ‘chelicerate + myriapod’ clade, but not basal arthropods. Certain clades derived from morphological taxonomy, such as Mandibulata, Atelocerata, Schizoramia, Maxillopoda and Cycloneuralia, are inconsistent with these rRNA data. The 28S gene contained more signal than the 18S gene, and contributed to the improved phylogenetic resolution. Our findings are similar to those obtained from mitochondrial and nuclear (e.g., elongation factor, RNA polymerase, Hox) protein-encoding genes, and should revive interest in using rRNA genes to study arthropod and ecdysozoan relationships.

© 2003 Elsevier Inc. All rights reserved.

Keywords: Large-subunit ribosomal RNA; 28S rRNA; Small-subunit ribosomal RNA; Phylogeny; Protostome; Ecdysozoa; Arthropod

1. Introduction

The Ecdysozoa concept says that all phyla of animals that grow by molting a cuticular exoskeleton are united in a monophyletic clade (Ecdysozoa = Arthropoda, Tardigrada, Onychophora, Nematoda, Nematomorpha, Priapulida, Kinorhyncha, and Loricifera). Originally derived from a phylogenetic study using 18S rRNA

genes (Aguinaldo et al., 1997), this controversial concept has been supported by subsequent molecular studies based on several genes (de Rosa et al., 1999; Giribet, 2002; Manuel et al., 2000; but see Hausdorf, 2000). However, many morphologists do not accept it (Brusca et al., 2003; Nielsen, 2001; Scholtz, 2002; Wägele and Misof, 2001; but see Peterson and Eernisse, 2001). We accept the Ecdysozoa because we recently obtained strong support for it from 28S rRNA genes (Mallatt and Winchell, 2002). The present paper explores the relationships among the different ecdysozoan lineages, which have also generated controversy (Giribet et al., 2000; Schmidt-Rhaesa et al., 1998). For example, some

[☆] Supplementary data associated with this article can be found at doi:10.1016/S1055-7903(03)00290-2.

* Corresponding author. Fax: +509-335-3184.

E-mail address: jmallatt@mail.wsu.edu (J.M. Mallatt).

morphological studies unite priapulans, kinorhynchs, loriciferans, nematodes, and nematomorphs into a monophyletic Cycloneuralia (Schmidt-Rhaesa et al., 1998) or Introverta (Nielsen, 1995), but some 18S-based studies indicate this group is paraphyletic (Garey, 2001; Peterson and Eernisse, 2001). Other controversies involve the relationships within Panarthropoda (= Arthropoda + Onychophora + Tardigrada) (Dewel and Dewel, 1997; Giribet, 2002; Peterson and Eernisse, 2001), and especially, relationships within Arthropoda. Fig. 1 shows four major hypotheses of the interrelationships of the arthropod subgroups: hexapods, myriapods, crustaceans, and chelicerates (also see Giribet and Ribera, 2000). Finally, the position of pycnogonids within arthropods is uncertain (Blaxter, 2001).

The goal of this study was to sequence nearly complete 28S and 18S rRNA genes from a large sample of ecdysozoans in order to resolve their relationships. This was pursued because of encouraging preliminary results (Mallatt and Winchell, 2002), and because rRNA seems to contain more signal than other genes used for high-order animal phylogeny (Giribet, 2002).

Analysis of rRNA gene sequences from 35 ecdysozoan and eight outgroup taxa using likelihood-based Bayesian inference and minimum evolution of LogDet-transformed distances provided important phylogenetic insights. These include the recovery of a basal position for Priapulida (and presumably for the related Kinorhyncha and Loricifera), the monophyly and internal relationships of Nematoda, paraphyly of Cycloneuralia, the monophyly of Panarthropoda and of Arthropoda, and a sister-group relationship between Pancrustacea

(= Crustacea + Hexapoda) and a lineage comprising Chelicerata and Myriapoda. These results emphasize the value of adding 28S sequences to 18S sequences for phylogenetic reconstruction and suggest that purported weaknesses of rRNA genes are attributable to inadequate sequence lengths and sub-optimal methods of phylogeny reconstruction used in the past. Bayesian inference performed extremely well on this data set, recovering established clades that are not obtained by other tree-building algorithms.

2. Materials and methods

2.1. Specimens and sequences

The names of the taxa sampled are shown in Figs. 2–4, whereas more detailed information, including voucher-specimen numbers and GenBank-accession numbers (AY210803-45), is presented in the Supplementary material (S1). Among the 35 ecdysozoans sampled, the large-subunit rRNA genes (28S + 5.8S) of 25 species were newly sequenced here. The taxa provide broad representation within Ecdysozoa, and comprise two priapulans, two nematomorphs, five nematodes, one tardigrade, two onychophorans, five chelicerates, four myriapods, eight crustaceans, and six hexapods. No kinorhynchs or loriciferans were available. Eight non-ecdysozoan bilaterians were used as outgroups, and provide a wide range of conserved lophotrochozoan and deuterostome sequences. DNA extraction, primers, PCR amplification, DNA purification, and sequence and fragment assembly followed protocols described in past studies (Mallatt and Sullivan, 1998; Winchell et al., 2002). The gene sequences were imported into SeqLab (Smith et al., 1994). There, the concatenated 28S and 18S genes were aligned by eye, with the alignment rigidly based on the large-subunit rRNA secondary-structure model of the frog *Xenopus laevis* (Schnare et al., 1996) and the small-subunit models of *X. laevis* and the sea urchin *Strongylocentrotus purpuratus* (Gutell, 1994). For the phylogenetic analyses, we used only the 28S and 18S genes, excluding the small 5.8S gene because it could not be sequenced across all ecdysozoan groups.

The 28S gene has a conserved *core* and 12 variable *divergent domains* (Hassouna et al., 1984; Hillis and Dixon, 1991; Mallatt et al., 2001). As in our past studies of deep-level phylogeny, we excluded the divergent domains and used only the core. For 18S, about 13% of all sites in the gene could not be aligned and were excluded. Overall, in the analyses we used 2313 and 1544 aligned sites from the 28S and 18S genes, respectively, for a total of 3857 sites. The alignment is available at <http://chuma.cas.usf.edu/~garey/alignments/alignment.html>.




Fig. 1. Major hypotheses of the relationships of arthropod groups: (A) classical 'Mandibulata' (= Mandibulata + Chelicerata), in which the three groups possessing mandibles formed by the second post-oral appendage (hexapods, myriapods, crustaceans) are distinguished from chelicerates; (B) Atelocerata versus Schizoramia: clades with unbranched versus two-branched appendages; (C) Pancrustacea + Myriapods within Mandibulata: that is, in the Mandibulata, hexapods group with crustaceans instead of with myriapods; (D) 'Chelicerata + Myriapoda' versus 'Crustacea + Hexapoda.'




Fig. 2. Bayesian inference, 50% consensus trees from combined 28S and 18S rRNA genes. The tree at left (A) is for all 43 taxa, a data set with nonstationary nucleotide frequencies and some divergent sequences. At right (B) is the tree for 37 taxa whose sequences exhibit stationary nucleotide frequencies. The numbers at the nodes are posterior probabilities, expressed as percentages. Asterisks indicate taxa whose 28S genes were not sequenced in this laboratory (all others were: see Supplementary material, S1).

2.2. Phylogenetic analyses

Phylogenetic algorithms that model the processes of sequence evolution, such as nucleotide substitution rates and among-site rate heterogeneity, are often preferred over algorithms that are not explicitly model-based (Bollback, 2002; Douady et al., 2003; Huelsenbeck et al., 2001b; Sullivan and Swofford, 2001). Here, we used two model-based algorithms for tree reconstruction: minimum evolution using LogDet-Paralinear distances (ME-LogDet) and Bayesian inference based on the likelihood function. These algorithms were executed using PAUP* 4.0 beta 10 (Swofford, 2001) or MrBayes 2.01 (Huelsenbeck and Ronquist, 2001).

LogDet is a model-based distance method for estimating phylogenetic trees (Lake, 1994; Lockhart et al., 1994; Swofford et al., 1996) that can be used with the ME algorithm. The ME-LogDet method models nucleotide-substitution rates better than other methods do and is designed to perform best when nucleotide frequencies are not stationary across taxa. By contrast, LogDet distances

do not account for among-site rate heterogeneity. PAUP* crudely handles such heterogeneity by dividing the gene into two parts: a proportion of sites that cannot vary, called P_i ; and all other sites, which are assumed to have evolved at the same rate. Following our previous studies, we chose a value of P_i (0.45) slightly lower than the proportion of sites that was constant across all 43 taxa (~0.5) (see Mallatt and Winchell, 2002; Winchell et al., 2002). This choice should be acceptable because trial analyses showed that all P_i values from 0.3–0.5 produced the same tree topology and similar bootstrap-support values. After the P_i was chosen, optimal ME-LogDet trees were calculated, and nonparametric bootstrap analyses were performed with 1000 replicates.

Bayesian inference based on the likelihood function (Huelsenbeck et al., 2001a,b; Larget and Simon, 1999) was used instead of the better-known maximum likelihood procedure (ML: Swofford et al., 1996) because it computes more efficiently and could analyze our 43 sequences in a reasonable amount of time (Hall, 2001; Huelsenbeck et al., 2002; Leache and Reeder, 2002).




Fig. 3. ME-LogDet distance trees (combined 28S and 18S genes): (A) optimal 43 taxon tree; (B) optimal 37 taxon tree. $P_i = 0.45$. At the nodes, ME-LogDet bootstrap percentages ($>50\%$, with some exceptions) are written above MP bootstrap percentages. Note that in the optimal LogDet tree in (A), *Hanseniella* goes with *Milnesium*; however, in the bootstrap consensus trees, *Hanseniella* goes with the onychophorans *Peripatoides* and *Peripatus* with strong bootstrap support of at least 95%.

However, it is more susceptible than ML to the error of assigning high confidence to short, incorrect nodes (Alfaro et al., 2003; Suzuki et al., 2002), so we used ML on subsets of sequences to check some of the Bayesian results (see below).

When running Bayesian analyses in MrBayes 2.01, we specified a general-time-reversible (GTR) model of nucleotide substitution with a proportion of invariable sites (I) and a gamma (Γ) distribution of among-site rate heterogeneity with four rate categories (see Mallatt and Sullivan, 1998; and Mallatt and Winchell, 2002). No initial values were assigned to the model parameters, and empirical nucleotide frequencies were used. Four Markov chains were run for a million generations and sampled every 100 generations to yield a posterior probability distribution of 10,000 trees. After eliminating the first 2000 trees as “burn-in,” we constructed a 50% majority-rule consensus tree, with nodal values representing the probability (“posterior probability”) that the recovered clades exist, given the aligned sequence data. We accepted a clade in the Bayesian tree at around 95% posterior probability (Murphy et al., 2001; Wilcox et al., 2002), while accepting values around 60–70% in the nonparametric bootstrap trees (Hillis and Bull, 1993).

A weakness of Bayesian inference, and of other algorithms based on the likelihood function, is that they assume stationarity of nucleotide frequency across all taxa in the data set, so their trees can be inaccurate if nonstationarity exists (Omilian and Taylor, 2001).

Finally, for comparison with past studies, we constructed trees using unweighted maximum parsimony (MP) (Swofford et al., 1996), including nonparametric bootstrapping with 1000 replicates. MP is not explicitly model-based and its resolving capacity may be compromised by a high susceptibility to long-branch attraction artifacts (Felsenstein, 1978; Lewis, 2001; Swofford et al., 2001).

2.3. Statistical tests of alternate hypotheses

Parametric bootstrapping based on ML (Efron, 1985; Huelsenbeck et al., 1996) was used to test if previous, alternate hypotheses of arthropod relations are consistent with present findings. Parametric bootstrapping can be extremely powerful and makes fewer simplifying assumptions than do other techniques of hypothesis testing. However, it is heavily reliant on the accuracy of the evolutionary model it uses to simulate sequence data




Fig. 4. Bayesian trees calculated from 28S (A) and from 18S (B) rRNA sequences. Forty-three taxa, 50% consensus trees. D1–D12 indicate the ecdysozoan nodes where the two trees differ, as listed and explained in Section 3.3.

(Goldman et al., 2000) and it may reject hypotheses too readily (Antezana, 2003).

To accommodate the high computational demands of ML-based parametric bootstrapping, we trimmed the data set to 26 conserved sequences by removing divergent sequences. Care was taken to assure that the retained sequences represented all major ecdysozoan groups and yielded a topology similar to that of the 43-taxon tree. Initially, the optimal, unconstrained, ML tree was estimated from the 26-taxon data, using an iterative search strategy (Sullivan et al., 1997), in which the ‘GTR + I + Γ ’ model was found to fit the data best. After this, the actual parametric bootstrap tests were performed, involving 100 simulations per hypothesis tested, as described elsewhere (Mallatt and Sullivan, 1998; Wilcox et al., 2002). Data were simulated with the Seq-Gen v1.2.3 program (Rambaut and Grassly, 2001).

3. Results

3.1. Unusual sequences and nucleotide frequencies

As the 43 gene sequences were being aligned, two things became evident: The rRNA genes of the two

nematomorphs, *Gordius* and *Chordodes*, are extremely similar, and those of the symphylian myriapod, *Hanseniella*, are highly divergent. The similarity between the two nematomorphs was so high (>98% for their entire 28S genes, 95% for their entire 18S genes) that it is doubtful whether they can be considered two distinct taxa in the phylogenetic analysis. The *Hanseniella* sequence is so different from all others that the first 350 nucleotides at the 5' end of its 18S gene were almost not alignable with the other sequences (but the rest of this 18S gene, and the 28S gene, were alignable); this divergence is reflected in the long branches for *Hanseniella* in the subsequent phylogenetic trees.¹

¹ The 18S sequence we obtained for *Hanseniella* was >1970 nucleotides long, whereas that obtained by Giribet and Wheeler (2001), from the same *Hanseniella* specimens, was only 1350 nucleotides. We cannot fully explain this discrepancy, but stand by our results. We obtained this sequence, without difficulty, by our usual method of amplifying the entire gene with two universal end-primers for 18S (5' primer: CTGGTTGATCCTGCCAGT; 3' primer: TA-ATGATCCTTCCGCAGGTTACCT). Furthermore, the shared parts of our and Giribet and Wheeler's sequences (AY210823 and AF173237) are fully identical in nucleotide sequence. Perhaps Giribet and Wheeler, who amplify their 18S genes in several pieces instead of one, did not amplify the ultra-divergent 5' end in this case?

When the χ^2 test of stationarity of nucleotide frequencies was applied to the 28S+18S data, the frequencies were found to be nonstationary across the 43 taxa ($\chi^2 = 336.5$; $P < 0.000000005$: see Supplementary material, S2). The 18S sequences were stationary, but the 28S were not, so the nonstationarity must arise from the 28S genes.

Given the nonstationarity of the entire data set, we identified the largest subset of sequences that was both stationary for nucleotide frequencies and biologically informative. To this end, we first removed *Chordodes* because it was nearly identical to *Gordius*. Then we removed the five divergent sequences with the most atypical proportions of nucleotides (Supplementary material, S2): C-rich *Hanseniella*, AT-rich *Drosophila* and *Aedes*; T-rich *Caenorhabditis*, and CG-rich *Peripatus*. The remaining 37 sequences were stationary ($\chi^2 = 124$; $P = 0.139$).

3.2. Combined 28S+18S trees

Figs. 2 and 3 show the Bayesian and ME-LogDet trees, respectively, for all 43 taxa and also for the stationary, 37-taxon subset. The 37-taxon trees are included because they best fit the assumptions of both the Bayesian method (for stationarity) and the ME-LogDet method (for the fewest divergent sequences likely to have experienced extreme evolutionary-rate heterogeneity among sites). The 43-taxon set is included because it is more complete, even though it violates stationarity and contains divergent “long-branched” sequences from *Hanseniella*, *Drosophila*, *Aedes*, *Caenorhabditis*, and *Peripatus* (branch lengths are indicated in Fig. 3 and in Supplementary material, S3).

Although not identical, the four trees in Figs. 2 and 3 share many similarities:

1. Priapulans are basal to the other Ecdysozoa, and Cycloneuralia (priapulans, nematomorphs, and nematodes) are paraphyletic.
2. All nematodes go together, with one subgroup consisting of *Trichinella* and *Xiphinema* and the other

containing *Ascaris* and *Meloidogyne* (and *Caenorhabditis*).

3. The nematodes go with the nematomorphs, although support for this ‘Nematoida’ node did not reach the level of significance in both Bayesian trees (e.g., 79% in Fig. 2B).
4. All panarthropods go together.
5. Chelicerates go with millipede and centipede myriapods.
6. The euchelicerates go together (scorpion *Pandinus*, xiphosuran *Limulus*, spider *Aphonopelma*).
7. Pycnogonids, *Callipallene* and *Colossendeis*, are in the ‘chelicerate + myriapod’ clade.
8. The Pancrustacea (hexapods and crustaceans) are united.
9. In Pancrustacea, the branchiopods go together (*Artemia*, *Daphnia*, *Triops*), as do the malacostracans (*Squilla* and *Panulirus*), but the maxillopods do not (*Argulus*, *Cyprididae* sp., *Cyclopidae* sp.).
10. The collembolan (*Podura*) goes with other hexapods.

On the other hand, there are several differences between the Bayesian and ME-LogDet trees, such as slightly different placements of pycnogonids within the ‘chelicerate + myriapod’ clade and different arrangements of traditional crustacean taxa in Pancrustacea. The biggest difference, however, is that ME-LogDet puts the long-branched sequences of the myriapod *Hanseniella*, and of the hexapods *Drosophila* and *Aedes*, outside the myriapod and hexapod clades, respectively (Fig. 3A), but Bayesian inference puts these sequences in their traditional clades (Fig. 2A). We checked the Bayesian result by individually adding *Hanseniella* and *Drosophila* to a 26-taxon subset of conserved sequences that was small enough for ML to handle (taxa listed at end of Table 1), and found that ML likewise put these two divergent sequences in their traditional clades.

3.3. 28S tree versus 18S tree

This section compares a tree derived from the 28S sequences alone with a tree from the 18S sequences

Table 1
Results of hypothesis testing by parametric bootstrapping

Alternate hypothesis	δ value	Range of 100 simulated difference values	P	Reject or accept?
1. Mandibulata consisting of Pancrustacea + Myriapoda (Fig. 1C)	47.5	0–6.34	$P \ll 0.01$	Reject
2. No ‘chelicerates + myriapods’ clade	12.0	0–5.07	$P \ll 0.01$	Reject
3. Pycnogonids as basal arthropods	7.8	0–3.76	$P \ll 0.01$	Reject
4. Pycnogonids as basal chelicerates	1.3	0–2.71	$P = 0.06$	Accept

Note. The test statistic, δ , is the difference between the ln likelihood value of the optimal tree (–27717.57) and the tree constrained to fit the alternate hypothesis (see Mallatt and Sullivan, 1998). The following 26 combined 28S+18S rRNA sequences were used (also see Supplementary material, S6): *Amphiporus*, *Aphonopelma*, *Argulus*, *Ascaris*, *Callipallene*, *Cormocephalus*, *Ctenolepisma*, *Cyclopidae*, *Cyprididae*, *Gordius*, *Gromphadorhina*, *Halicryptus*, *Limulus*, *Lithobius*, *Milnesium*, *Orthoporus*, *Pandinus*, *Peripatoides*, *Placopecten*, *Podura*, *Ptychodera*, *Squilla*, *Stylochus*, *Tenebrio*, *Triops*, *Xiphinema*.

(see Fig. 4, with their branch lengths shown in Supplementary material, S4–S5). Because Bayesian inference seems to place long-branched sequences more accurately than does ME-LogDet (see above), we show only the Bayesian trees here. In this section, tree topology is considered more important than high nodal support because the goal is to find which of the two gene trees most closely resembles the combined 28S+18S tree. Both the 28S (Fig. 4A) and 18S (Fig. 4B) trees are roughly similar to the combined-gene tree (Fig. 2A) and to one another, sharing many nodes. For example, both 28S and 18S trees show strong support for priapulans as basal ecdysozoans, for paraphyly of Cycloneuralia, for two groups of nematodes (*Trichinella* and *Xiphinema* versus *Ascaris*, *Meloidogyne* and *Caenorhabditis*), and for monophyletic euchelicerates, Pancrustacea, branchiopods, and malacostracans. On the other hand, the 28S and 18S trees differ in the following ecdysozoan clades (Fig. 4: D1–D12):

- D1. *Nematomorphs*. In the 28S tree, the nematomorphs (*Chordodes* and *Gordius*) are basal to nematodes and panarthropods, but in the 18S tree they group with the nematodes.
- D2. *Nematodes*. Within the nematodes, the relative positions of *Ascaris*, *Meloidogyne*, and *Caenorhabditis* differ.
- D3. *Strong Panarthropoda*. The 28S tree has more support for Panarthropoda (100% versus 61%).
- D4. *Pycnogonids*. The 28S tree places the pycnogonids (*Callipallene* and *Colossendeis*) in the ‘chelicerate + myriapod’ clade, but the 18S tree shows them as distinct from euchelicerates and from a ‘myriapod + pancrustacean’ line.
- D5. *Myriapods*. The 28S tree groups myriapods with chelicerates, but the 18S tree places centipede and millipede myriapods as the sister to Pancrustacea.
- D6. *Hanseniella*’s 28S sequence goes with the myriapods, but its 18S sequence forms a distinct lineage in Panarthropoda.
- D7. In crustaceans, 28S places *Argulus* at the base of the Pancrustacea, whereas 18S puts the branchiopods in this basal position.
- D8. In branchiopods, 28S joins *Triops* with *Daphnia*, but 18S joins *Triops* with *Artemia*.
- D9. *Cyclopidae sp.* 28S groups this copepod with hexapods, 18S puts it with malacostracans.
- D10. *Podura*. The 28S sequence of *Podura* groups with the other hexapod sequences, but its 18S sequence falls outside the hexapods.
- D11. 28S fails to unite *Gromphadorhina* and *Tenebrio*, but 18S does unite these pterygote insects.
- D12. *Dipteran hexapods*. Although neither gene correctly places the dipterans *Drosophila* and *Aedes* with the other pterygote insects *Gromphadorhina* and *Tenebrio*, the 18S gene joins them with silverfish *Ctenolepisma*, an apterygote.

The above list indicates that the 28S gene has the stronger influence on the combined 28S + 18S tree. That is, for eight of the 12 differences (D3–D10), the 28S condition is seen in the combined tree of Fig. 2A. By contrast, the 18S condition dominates in only four cases (D1–D2, D11–D12).

3.4. Maximum parsimony tree

Maximum-parsimony analysis of the 28S + 18S sequences produced trees that were more like the ME-LogDet trees than the Bayesian trees, so parsimony-bootstrap values are included on the ME-LogDet trees in Fig. 3. The MP bootstrap-consensus trees (not shown) resembled the ME trees in misplacing the divergent *Aedes*, *Drosophila* and *Hanseniella* sequences, but were unique in that they did not unite chelicerates with myriapods, nor centipedes with the millipede.

3.5. Hypothesis testing

Parametric-bootstrapping of a 26-taxon subset of the 28S + 18S sequences tested four alternative hypotheses of arthropod interrelationships (Table 1). As planned, the 26 sequences in this subset showed stationarity of nucleotide frequencies ($\chi^2 = 68.51$; $P = 0.69$) and they produced an ML tree (shown in Supplementary material, S6) that is similar to the Bayesian tree derived from the larger, stationary set of 37 sequences (Fig. 2B) in that it recovers the same major groups of arthropods. Three of the alternate hypotheses tested in Table 1 have received support from a whole-evidence study (Giribet et al., 2001): *hypothesis 1*. The basic groups of arthropods are Mandibulata and Chelicerata, with Mandibulata consisting of Pancrustacea and Myriapoda (Fig. 1C); *hypothesis 2*. Chelicerates + myriapods do not form a clade; *hypothesis 3*. Pycnogonids are not chelicerates as traditionally claimed, but instead are basal arthropods. Finally, we tested the traditional hypothesis 4 that pycnogonids are basal chelicerates (Waloszek and Dunlop, 2002). Results show that our rRNA data are consistent only with #4, pycnogonids as basal chelicerates ($\delta = 1.3$, $P = 0.06$), and are highly inconsistent with the other three hypotheses (in which δ values were > 7.8 and P was $\ll \ll 0.01$). The time-consuming computations of parametric bootstrapping kept us from using it to test other, more dissimilar, hypotheses of arthropod relations. However, we found that both Classical Mandibulata (Fig. 1A: $\delta = 184$) and Atelocerata + Schizoramia (Fig. 1B: $\delta = 202$) were strongly rejected ($P < 0.01$) by the SH test, which is far more conservative than parametric bootstrapping (Shimodaira and Hasegawa, 1999; Goldman et al., 2000).

4. Discussion

4.1. General comparison to previous studies

Most previous analyses of ecdysozoan phylogeny using 18S genes, with or without a small part of 28S, used MP or the parsimony-compatible ‘direct optimization method’ (Giribet and Ribera, 2000; Peterson and Eernisse, 2001; Spears and Abele, 1997; Wheeler, 1997). These analyses often yielded low resolution or questionable results, such as paraphyly of established clades and terminal taxa in unlikely positions, including failure to show the monophyly of Panarthropoda or Arthropoda, recovery of onychophorans and tardigrades within other ecdysozoan phyla, paraphyletic myriapoda in which centipedes did not unite with millipedes, placement of collembolans outside Hexapoda, placement of highly divergent sequences (e.g., symphylans, dipterans) outside their traditional groups, and uncertain and variable placement of the nematode clade (the latter reviewed by Garey, 2001). Consequently, some authorities expressed discouragement at the performance of rRNA genes (e.g., Giribet and Ribera, 2000; Spears and Abele, 1997), especially when the situation was not improved by adding more 18S sequences. Confirming these shortcomings, the 18S MP tree for our own 43 taxa also showed low resolution (Supplementary material, S7).

The shortcomings noted above have been largely eliminated from our combined 28S + 18S analysis. Although our taxon sampling remains low relative to the diversity (even hyper-diversity) of the represented groups, the present Bayesian trees are better resolved (Fig. 2) than trees from earlier studies and show little paraphyly of accepted clades. Specifically, our analyses recovered monophyletic Panarthropoda, Arthropoda, Myriapoda and Hexapoda, and a monophyletic Nematoda that fit cleanly in the Ecdysozoa. True resolving power is also suggested by the fact that ours is the only analysis to place the “most abnormal” arthropod rRNA sequence known (Giribet and Ribera, 2000)—that of a symphylan—into the myriapod clade, a placement that is supported by morphology and other genes (Edgecombe and Giribet, 2002; Giribet et al., 2001; Regier and Shultz, 2001). We attribute the improved resolution of the present study to three factors, namely, adding 28S to the 18S sequences, using the likelihood-based Bayesian method, and using a large enough sample of ecdysozoans (35).

These factors minimized long-branch attraction and other artifacts, but our method and results are not perfect. For example, in the outgroup, the chordates *Ciona* and *Hydrolagus* are not united (the difficulties in recovering a monophyletic Chordata from rRNA data were discussed by Winchell et al., 2002). As another example, the Bayesian tree (Fig. 2A) recovers erroneous relationships within Hexapoda, in that it places the

dipterans *Drosophila* and *Aedes* with silverfish *Ctenolepisma* instead of with the other holometabolous, pterygote insect *Tenebrio*. However, dipteran rRNA sequences are notoriously divergent, AT-biased, and likely to disrupt gene trees (Friedrich and Tautz, 1995; Omilian and Taylor, 2001). Bayesian inference at least placed the dipterans within Hexapoda, whereas ME-LogDet and MP incorrectly united them with branchiopod crustaceans (Fig. 2A versus Fig. 3A).²

Unlike most previous workers, Garey (2001) eschewed parsimony methods and used ME-LogDet analysis of 18S to investigate the relationships among ecdysozoan phyla (except loriciferans). Working before Bayesian methods or complete 28S sequences were available, he increased resolution by using only those taxa with the most-slowly evolving 18S sequences. The relationships he discerned were: ((Priapulida + Kinorhyncha), ((Nematoda + Nematomorpha), (Tardigrada, (Onychophora, (Arthropoda))))), with all nodes supported by bootstrap values of $\geq 69\%$, except that the relative positions of the Onychophora and Tardigrada were uncertain. This is fully consistent with the findings of the present 28S + 18S study (Fig. 2).

4.2. Comparison of ME-LogDet and Bayesian methods

Ecdysozoan sequences were analyzed here using both ME-LogDet and Bayesian inference based on the likelihood function (GTR + I + Γ model). Both of these methods have acknowledged weaknesses (see Section 2), and initially it was unknown which one would produce the most accurate trees from our data. We conclude that Bayesian inference is preferable because it placed the divergent dipteran and symphylan sequences in more reasonable positions than did ME-LogDet. Thus, for our data set, the strength of Bayesian inference in accounting for rate heterogeneity among sites in a gene seems to overshadow any weakness caused by nucleotide nonstationarity and overconfidence of the method. ME-LogDet misplaced the divergent sequences, apparently due to its limited ability to model rate heterogeneity. However, a test of this interpretation using bootstrapper’s gambit, a LogDet method that does model rate

² Immediately after this paper was accepted for publication, we solved this dipteran problem. The complete rRNA sequence of a lepidopteran (moth: Wang et al., 2003), widely held to represent the sister group of dipterans (Wheeler et al., 2001), became available, and we scoured GenBank to find nearly complete rRNA sequences from some additional pterygote insects: a hymenopteran (ant: Chalwatzis, 1995; Ohnishi, 2000), a hemipteran (alfalfa hopper: Campbell et al., 1995; Dietrich et al., 2001), and another dipteran (midge: Koepf et al., 1996). With these new sequences, both Bayesian analysis and ML placed the dipterans as sister to the moth in a monophyletic Pterygota, as predicted by conventional taxonomy (Wheeler et al., 2001), whereas ME-LogDet and MP continued to place the dipterans basal to all other hexapod sequences. The ML tree is presented in Supplemental material (S8).

heterogeneity (Lake, 1998), yielded mixed results: Gambit correctly put dipterans in insects but still incorrectly joined the ultra-divergent *Hanseniella* with onychophorans (results not shown).

It should be noted that in our earlier studies, which examined much higher-order relationships among animals, likelihood-based methods seemed to provide *worse* phylogenetic resolution than the ME-LogDet method (Mallatt and Winchell, 2002; Winchell et al., 2002), which is the opposite of what we found here within the Ecdysozoa. This situation may reflect the sparser taxon sampling of the earlier studies (often just one rRNA sequence per phylum), which may have provided too little information to the likelihood algorithm.

4.3. Comparison of 28S and 18S trees

The trees derived from the separate analyses of 28S and 18S sequences have roughly similar topologies (Fig. 4), but we regard the 28S gene as contributing more phylogenetic signal. Specifically, 28S alone puts the collembolan *Podura* with the other hexapods and also puts the divergent *Hanseniella* and dipterans in more reasonable places (compare Figs. 4A and 4B). Also, as explained in Section 3.3, the 28S gene has more influence on the topology of the combined-gene tree than does the 18S gene.

These relative contributions of 28S versus 18S are consistent with the findings of previous studies. Most prior 28S + 18S studies dealt with higher-order clades than used here, namely metazoans and bilaterians at the “phylum”-to-“superphylum” level, and found 18S to contribute more signal than 28S (Mallatt and Winchell, 2002; Medina et al., 2001; Winchell et al., 2002). By contrast, a study of lower-level phylogeny, within the “class” of cartilaginous fishes, found that nearly all the signal came from the 28S gene (Winchell, 2001). The present study covered an intermediate taxonomic level, from “phyla” to “classes” within Ecdysozoa, and found 28S to provide just slightly more signal than 18S. Thus, a trend is evident: In proceeding from high to low taxonomic levels, the 28S gene contributes more signal and the 18S gene less.

4.4. Comparison of clade-specific findings with those of previous studies

4.4.1. Cycloneuralia

The clade Cycloneuralia (Ehlers et al., 1996; Schmidt-Rhaesa et al., 1998) or Introverta (Nielsen, 1995) was proposed to include Scalidophora (Priapulida + Kinorhyncha + Loricifera) and Nematoida (Nematoda + Nematomorpha). The morphological characters used to unite this clade were a circumpharyngeal cerebral ring that is not ganglionated anterodorsally, details of brain histology, and perhaps an introvert organ. Our

findings are inconsistent with the Cycloneuralia concept because they place priapulans basal to all other ecdysozoans (Figs. 2 and 3). The other scalidophorans were not sampled here but should be united with priapulans: Indeed, in his analysis of conserved 18S sequences, Garey (2001) recovered kinorhynchs and priapulans as the basal, monophyletic group in Ecdysozoa. If Cycloneuralia are paraphyletic, their shared anatomical characters must either be primitive or convergently evolved.

4.4.2. Nematoida

Schmidt-Rhaesa (1998) identified morphological characters that appear to unite the nematodes and nematomorphs into the taxon Nematoida. These synapomorphies include aflagellate sperm, a cloaca in both sexes, and similarities in the ventral and dorsal epidermal and nerve cords. Most 18S-based studies have not recovered Nematoida (Giribet et al., 2000; Peterson and Eernisse, 2001), but it was supported in Garey’s (2001) analysis of conserved 18S sequences. The present analysis of 28S + 18S sequences also recovered a Nematoida, but this support was weak in that it lost significance when one of the two nearly identical nematomorph sequences was omitted (Fig. 2B). Future studies should include the other nematomorph clade, the presumably primitive marine *Nectonema* (Bleidorn et al., 2002; Schmidt-Rhaesa, 2002).

4.4.3. Relationships among nematodes

We used only five species of nematodes, so we cannot say much about relationships within this large phylum. As seen in Figs. 2 and 3, our trees show two groups, *Xiphinema* + *Trichinella* versus *Ascaris* + *Caenorhabditis* + *Meloidogyne*, with the relationships within the latter group unresolved. These two groups are consistent both with morphology-based systematics of these genera (adenophoreans versus secernenteans: Blaxter et al., 1998, their Fig. 2b) and with a tree based on 18S (Blaxter et al., 1998, their Fig. 2a).

4.4.4. Panarthropoda

A panarthropod clade consisting of onychophorans, tardigrades and arthropods is widely accepted, based on such shared characters as a segmental body and ventrolateral appendages (Schmidt-Rhaesa et al., 1998). Although this clade was never well supported by 18S, it was strongly supported by 28S + 18S here. However, the relations among the three panarthropodan phyla are uncertain (Dewel and Dewel, 1997; Peterson and Eernisse, 2001), and all possible pairings have been advocated (Eriksson et al., 2003; Giribet, 2002; Ramsköld and Chen, 1998; Schmidt-Rhaesa et al., 1998). Our findings are also inconclusive, but hint at ‘onychophorans + tardigrade’ (Fig. 2). Adding more tardigrade sequences might clarify these relations, but the divergent

onychophoran sequences might confound attempts to improve resolution.

4.4.5. *Arthropod subphyla*

The phylogenetic relationships of the major arthropod groups (Chelicerata, Crustacea, Myriapoda and Hexapoda) are intensely debated. Prior to recent molecular studies, most neontologists favored ‘Mandibulata + Chelicerata’ (Fig. 1A), which unites hexapods, myriapods, and crustaceans to the exclusion of chelicerates (Snodgrass, 1938). By contrast, most paleontologists favor Atelocerata (= Hexapoda + Myriapoda) and Schizoramia (= Chelicerata + Crustacea) (Fig. 1B) (Cisne, 1974; Colgan et al., 1998; Haas et al., 2003; Wills et al., 1998). Both these hypotheses originally included the monophyly of ‘Hexapoda + Myriapoda,’ but molecular studies now strongly suggest that Hexapoda and Crustacea form a monophyletic group, the Pancrustacea (= Tetraconata: Dohle, 2001) (Giribet et al., 2001; Hwang et al., 2001; Regier and Shultz, 2001; Zrzavy and Stys, 1997; see Giribet and Ribera, 2000, p. 222 for other references). Correspondingly, certain morphological characters are shared by hexapods and crustaceans, especially in the nervous system (Mittmann, 2002; Richter, 2002; Wilson et al., 2000). Some studies suggest that Pancrustacea is the sister group to Myriapoda within Mandibulata (Fig. 1C) (Giribet et al., 2001), but others indicate that Pancrustacea is sister to a clade comprising Chelicerata and Myriapoda (Cook et al., 2001; Hwang et al., 2001). Our results support this ‘chelicerate + myriapod’ hypothesis (Figs. 1D and 2), and reject Mandibulata + Chelicerata and Atelocerata + Schizoramia. Even the Pancrustacea + Myriapoda version of Mandibulata (Fig. 1C), which is the least inconsistent with our trees, was overwhelmingly rejected by parametric bootstrapping (Table 1).

The hypothesis that chelicerates and myriapods form a monophyletic sister group to Pancrustacea was never supported by morphology-based studies and it derives strictly from molecular studies. Friedrich and Tautz (1995) were early supporters, based on their analysis of partial-28S/18S sequences from ten arthropods (also see Turbeville et al., 1991; and Giribet et al., 1996), but the best evidence for ‘chelicerates + myriapods’ has come from Hox- and mitochondrial gene sequences (Cook et al., 2001; Hwang et al., 2001). Given our new rRNA evidence that this clade might actually exist, we name it “Paradoxopoda,” which reflects the conflict between the molecular data that support it and the morphological data that do not.

Overall, our findings resemble those of Regier and Shultz (2001), who explored arthropod interrelationships with the nuclear genes for RNA polymerase II, elongation factor-1 α , and elongation factor 2. The number of taxa they used (19 panarthropods) and the number of nucleotide sites in their combined-gene se-

quences (~4000) are similar to what we used here (26 panarthropods, ~3800 nucleotides). As indicated by the ML-bootstrap values in their three-gene summary table (their p. 142), they found monophyly of arthropods (>87%); of myriapods including a symphylan (94–99%); of the chelicerates including pycnogonids (75–86%); of Pancrustacea (95–97%); and less strongly, of hexapods (including collembolans) (66–87%). Their only finding that differed from ours was that the ‘chelicerate + myriapod’ clade was never supported.

4.4.6. *Chelicerates*

Pycnogonids are usually considered chelicerates, but their position is controversial because they have many autapomorphic structures (Blaxter, 2001; Brusca and Brusca, 1990). However, pycnogonids primitively have chelicerae, a key synapomorphy of chelicerates, and Waloszek and Dunlop (2002) argued in detail that their body plan fits that of chelicerates. By contrast, Giribet et al. (2001) concluded that pycnogonids are at the base of all extant arthropods, based on a synthesis of characters from 18S, other genes, and morphology (this placement is also consistent with our 18S tree in Fig. 4B). Our combined 28S + 18S data indicate that pycnogonids are not basal arthropods but are either basal chelicerates or a distinct lineage in the ‘chelicerate + myriapod’ clade (Figs. 2 and 3; Table 1).

4.4.7. *Myriapoda*

Our findings favor the traditional view of myriapod monophyly (Edgecombe and Giribet, 2002; Giribet et al., 2001; Regier and Shultz, 2001; Wills et al., 1998) over myriapod paraphyly (Kraus, 2001; Shear, 1997) because our trees separate myriapods from hexapods (Fig. 2). Myriapods are the group that would benefit most from more rRNA sequences.

4.4.8. *Pancrustacea*

The monophyly of Pancrustacea was strongly supported by our 28S and 18S sequences (Figs. 2–4). Among the traditional crustaceans, our rRNA trees indicated that branchiopods (*Artemia*, *Daphnia*, and *Triops*) and eumalacostracans (*Squilla* and *Panulirus*) are each monophyletic, a result also indicated by 18S and other genes (Shultz and Regier, 2000; Spears and Abele, 1997). Maxillopoda, by contrast, is problematical. This clade, said to consist of copepods, ostracods, branchiurans, cirripeds, and mystacocarids, was united by such morphological characters as body segmentation and mouth parts (Waloszek and Müller, 1998; Wills et al., 1997). However, it was never universally accepted by systematists (see Spears and Abele, 1997) and was never supported by molecular studies (Giribet and Ribera, 2000; Regier and Shultz, 2001; Shultz and Regier, 2000), including the present study, in which the cyclopid

copepod, cypridid ostracod, and branchiuran *Argulus* do not group as a unit (Figs. 2–4).

Our study, like all previous rRNA-based studies, indicates that hexapods nest within a paraphyletic “crustacea,” which raises the question of which crustacean line is the sister group to Hexapoda (Friedrich and Tautz, 2001; Schram and Jenner, 2001). Although other studies favor either branchiopods (Gaunt and Miles, 2002; Regier and Shultz, 1997) or malacostracans (Hwang et al., 2001; Wilson et al., 2000), our Bayesian trees (Fig. 2) show a copepod as the sister of hexapods. However, the copepod sequence is rather divergent and copepods have never been regarded as relevant to understanding the origin of hexapods.

Finally, our 28S + 18S data unite the collembolan *Podura* with the other hexapods (Fig. 2), supporting the traditional idea of hexapod monophyly (cf. Nardi et al., 2003).

5. Conclusions

Overall, our findings suggest that recent pessimism about the ability of rRNA sequences to resolve ecdysozoan interrelationships can be replaced with optimism. The use of likelihood-based Bayesian inference on nearly complete 28S + 18S sequences may prove to be the long-awaited combination of best genes (Giribet, 2002) and best tree-building method for reconstructing ecdysozoan phylogenies with the fewest long-branch-attraction artifacts. This implies that adding more taxa will finally start to improve, rather than diminish, phylogenetic resolution. It is not meant to imply, however, that rRNA alone is sufficient. Instead, many genes must be used (Brown et al., 2001; Takezaki et al., 2003; Sanderson et al., 2003). To this end, the rRNA sequences could be concatenated with sequences of nuclear protein-coding and mitochondrial genes, which should produce even better taxonomic resolution.

6. Summary

In this study, the use of nearly complete 28S + 18S rRNA sequences and of Bayesian inference based on a GTR + I + Γ likelihood model produced a well-resolved tree of 35 arthropod and ecdysozoan sequences. More specifically, our trees support the following findings (Fig. 2).

1. Priapulans (and presumably the related kinorhynch and loriciferans) represent the basal line of ecdysozoans.
2. The five nematodes used here form two subgroups consistent with accepted clades.
3. Arthropoda and Panarthropoda are each monophyletic, although the interrelationships of the panar-

thropod phyla (tardigrades, onychophorans, and arthropods) remain uncertain.

4. Arthropoda contains two major divisions: Pancrustacea (Hexapoda in a paraphyletic “crustacea”) and the newly named Paradoxopoda (‘Chelicerata + Myriapoda’).
 5. Euchelicerates, Hexapoda, ectognathic insects, and Myriapoda are each monophyletic.
 6. Pycnogonids are either chelicerates, or part of the ‘chelicerate + myriapod’ clade.
 7. Mandibulata, Maxillopoda, Atelocerata, Schizoramia, and Cycloneuralia are paraphyletic and invalid taxa.
- Other major findings are:
1. In this data set, the 28S gene contains more phylogenetic signal than the 18S gene.
 2. In this study, minimum-evolution analysis of LogDet distances and maximum parsimony produced less resolution than Bayesian inference.
 3. Arthropod relationships derived here from rRNA genes resemble those calculated from concatenated elongation-factor and RNA polymerase II genes (Regier and Shultz, 2001).

Acknowledgments

Thanks are extended to Gary Thorgaard, Jack Sullivan, Kevin Pullen, and to all who provided specimens.

References

- Aguinaldo, A.M.A., Turbeville, J.M., Linford, L.S., Rivera, M.C., Garey, J.R., Raff, R.A., Lake, J.A., 1997. Evidence for a clade of nematodes, arthropods, and other moulting animals. *Nature* 387, 489–492.
- Alfaro, M.E., Zoller, S., Lutzoni, F., 2003. Bayes or bootstrap? A simulation study comparing the performance of Bayesian Markov Chain Monte Carlo sampling and bootstrapping in assessing phylogenetic confidence. *Mol. Biol. Evol.* 20, 255–266.
- Antezana, M., 2003. When being “most likely” is not enough: examining the performance of three uses of parametric bootstrapping in phylogenetics. *J. Mol. Evol.* 56, 198–222.
- Blaxter, M., 2001. Sum of the arthropod parts. *Nature* 413, 121–122.
- Blaxter, M.L., De Ley, P., Garey, J.R., Liu, L.X., Scheldeman, P., Vierstraete, A., Vanfleteren, J.R., Mackey, L.Y., Dorris, M., Frisse, L.M., Vida, J.T., Thomas, W.K., 1998. A molecular evolutionary framework for the phylum Nematoda. *Nature* 392, 71–75.
- Bleidorn, C., Schmidt-Rhaesa, A., Garey, J.R., 2002. Systematic relationships of nematomorpha based on molecular and morphological data. *Invert. Biol.* 121, 357–364.
- Bollback, J.P., 2002. Bayesian model adequacy and choice in phylogenetics. *Mol. Biol. Evol.* 19, 1171–1180.
- Brown, J.R., Douady, C.J., Italia, M.J., Marshall, W.E., Stanhope, M.J., 2001. Universal trees based on large combined protein sequence data sets. *Nat. Genet.* 28, 281–295.
- Brusca, R.C., Brusca, G.J., 1990. *Invertebrates*. Sinauer, Sunderland, MA.

- Brusca, R.C., Brusca, G.J., Burness, T.J., 2003. Invertebrates, second ed. Sinauer, Sunderland, MA.
- Campbell, B.C., Steffen-Campbell, J.D., Sorensen, J.T., Gill, R.J., 1995. Paraphyly of Homoptera and Auchenorrhyncha inferred from 18S rDNA nucleotide sequences. *Syst. Entom.* 20, 175–194.
- Chalwatzis, N., 1995. Unpublished submission to GenBank, Accession number X89492.
- Cisne, J.L., 1974. Trilobites and the origin of arthropods. *Science* 186, 13–18.
- Colgan, D.J., McLauchlan, A., Wilson, G.D.F., Livingston, S.P., Edgecombe, G.D., Macaranas, J., Cassis, G., Gray, M.R., 1998. Histone H3 and U2 snRNA DNA sequences and arthropod molecular evolution. *Austr. J. Zool.* 46, 419–437.
- Cook, C.E., Smith, M.L., Telford, M.J., Bastianello, A., Akam, M., 2001. Hox genes and the phylogeny of the arthropods. *Curr. Biol.* 11, 759–763.
- Dietrich, C.H., Rakitov, R.A., Holmes, J.L., Black, W.C., 2001. Phylogeny of the major lineages of Membracoidea (Insecta: Hemiptera: Cicadomorpha) based on 28S rDNA sequences. *Mol. Phylogenet. Evol.* 18, 293–305.
- de Rosa, R., Grenier, J.K., Andreeva, T., Cook, C.E., Adoutte, A., Akam, M., Carroll, S.B., Balavoine, G., 1999. Hox genes in brachiopods and priapulids and protostome evolution. *Nature* 399, 772–776.
- Dewel, R.A., Dewel, W.C., 1997. The place of tardigrades in arthropod evolution. In: Fortey, R.A., Thomas, R.H. (Eds.), *Arthropod Relationships*. Chapman & Hall, London, pp. 109–123.
- Dohle, W., 2001. Are the insects terrestrial crustaceans? A discussion of some new facts and arguments and the proposal of the proper name 'Tetraconata' for the monophyletic unit Crustacea + Hexapoda. *Ann. Soc. Entomol. Fr. (N.S.)* 37, 85–103.
- Douady, C.J., Delsuc, F., Boucher, Y., Doolittle, W.F., Douzery, E.J.P., 2003. Comparison of Bayesian and maximum likelihood bootstrap measures of phylogenetic reliability. *Mol. Biol. Evol.* 20, 248–254.
- Edgecombe, G.D., Giribet, G., 2002. Myriapod phylogeny and the relationships of Chilopoda. In: Bousquets, J.E., Morrone, J.J., Ponce Ulloa, H. (Eds.), *Biodiversidad, taxonomía y biogeografía de artrópodos de México: hacia una síntesis de su conocimiento*. Pressas de Ciencias. Universidad Nacional Autónoma de México, México D.F., pp. 143–168.
- Efron, B., 1985. Bootstrap confidence intervals for a class of parametric problems. *Biometrika* 72, 45–58.
- Ehlers, U., Alrichs, W., Lemburg, C., Schmidt-Rhaesa, A., 1996. Phylogenetic systematization of the Nematelminthes (Aschelminthes). *Verh. Dtsch. Zool. Ges.* 89.1, 8.
- Eriksson, B.J., Tait, N.N., Budd, G.E., 2003. Head development in the onychophoran *Euperipatoides kanangrensis* with particular reference to the central nervous system. *J. Morph.* 255, 1–23.
- Felsenstein, J., 1978. Cases in which parsimony and compatibility methods will be positively misleading. *Syst. Zool.* 27, 401–410.
- Friedrich, M., Tautz, D., 1995. Ribosomal DNA phylogeny of the major extant arthropod classes and the evolution of myriapods. *Nature* 376, 165–167.
- Friedrich, M., Tautz, D., 2001. Arthropod rDNA phylogeny revisited: a consistency analysis using Monte Carlo simulation. *Ann. Soc. Entomol. Fr. (N.S.)* 37, 21–40.
- Garey, J.R., 2001. Ecdysozoa: the relationship between Cycloneuralia and Panarthropoda. *Zool. Anz.* 240, 321–330.
- Gaunt, M.W., Miles, M.A., 2002. An insect molecular clock dates the origin of the insects and accords with palaeontological and biogeographic landmarks. *Mol. Biol. Evol.* 19, 748–761.
- Giribet, G., 2002. Current advances in the phylogenetic reconstruction of metazoan evolution, A new paradigm for the Cambrian explosion? *Mol. Phy. Evol.* 24, 345–357.
- Giribet, G., Carrunza, S., Baguña, J., Riutort, M., Ribera, C., 1996. First molecular evidence for the existence of a Tardigrada + Arthropoda clade. *Mol. Biol. Evol.* 13, 76–84.
- Giribet, G., Distel, D.L., Polz, M., Sterrer, W., Wheeler, W.C., 2000. Triploblastic relationships with emphasis on the acoelomates and the position of Gnathostomulida, Cycliophora, Plathelminthes, and Chaetognatha: A combined approach of 18S rDNA sequences and morphology. *Syst. Biol.* 49, 539–562.
- Giribet, G., Edgecombe, G.D., Wheeler, W.C., 2001. Arthropod phylogeny based on eight molecular loci and morphology. *Nature* 413, 157–161.
- Giribet, G., Ribera, C., 2000. A review of arthropod phylogeny: new data based on ribosomal DNA sequences and direct character optimization. *Cladistics* 16, 204–231.
- Giribet, G., Wheeler, W.C., 2001. Some unusual small-subunit ribosomal RNA sequences of metazoans. *Am. Mus. Novit.* 3337, 14.
- Goldman, N., Anderson, J.P., Rodrigo, A.G., 2000. Likelihood-based tests of topologies in phylogenetics. *Syst. Biol.* 49, 652–670.
- Gutell, R.R., 1994. Collection of small subunit (16S- and 16S-like) ribosomal RNA structures. *Nucl. Acids Res.* 22, 3502–3507.
- Haas, F., Waloszek, D., Hartenberger, R., 2003. *Devonohexapodus bocksbergensis*, a new marine hexapod from the Lower Devonian Hunsrück Slates, and the origin of Atelocerata and Hexapoda. *Org. Divers. Evol.* 3, 39–54.
- Hall, B.G., 2001. *Phylogenetic Trees Made Easy: A How-To Manual for Molecular Biologists*. Sinauer Associates Inc, Sunderland, MA.
- Hassouna, N., Michot, B., Bachelier, J.-P., 1984. The complete nucleotide sequence of mouse 28S rRNA gene: implications for the process of size increase of the large subunit rRNA in higher eukaryotes. *Nucl. Acids Res.* 12, 3563–3583.
- Hausdorf, B., 2000. Early evolution of the Bilateria. *Syst. Biol.* 49, 130–142.
- Hillis, D.M., Bull, J.J., 1993. An empirical test of bootstrapping as a method for addressing confidence in phylogenetic analysis. *Syst. Biol.* 42, 182–192.
- Hillis, D.M., Dixon, M.T., 1991. Ribosomal DNA: molecular evolution and phylogenetic inference. *Q. Rev. Biol.* 66, 411–453.
- Huelsenbeck, J.P., Hillis, D.M., Jones, R., 1996. Parametric bootstrapping in molecular phylogenetics: applications and performance. In: Ferraris, J.E., Palumbi, S.R. (Eds.), *Molecular Zoology: Advances, Strategies, and Protocols*. Wiley-Liss, New York, pp. 19–45.
- Huelsenbeck, J.P., Ronquist, F., 2001. MrBayes: Bayesian inference of phylogeny. Version 2.01. Distributed by the author. Department of Biology, University of Rochester.
- Huelsenbeck, J.P., Ronquist, F., Hall, B., 2001a. MrBayes: A program for the Bayesian inference of phylogeny. *Instruction Manual for Version 2.01*, 17 p.
- Huelsenbeck, J.P., Ronquist, F., Nielsen, R., Bollback, J.P., 2001b. Bayesian inference of phylogeny and its impact on evolutionary biology. *Science* 294, 2310–2314.
- Huelsenbeck, J.P., Larget, B., Miller, R.E., Ronquist, F., 2002. Potential applications and pitfalls of Bayesian inference of phylogeny. *Syst. Biol.* 51, 673–688.
- Hwang, U.W., Friedrich, M., Tautz, D., Park, C.J., Kim, W., 2001. Mitochondrial protein phylogeny joins myriapods with chelicerates. *Nature* 413, 154–157.
- Koepf, H., Hankeln, T., Schmidt, E.R., 1996. Organization and evolution of the rDNA genes in Chironomus. Unpublished submission to GenBank, Accession number X99212.
- Kraus, O., 2001. "Myriapoda" and the ancestry of the Hexapoda. *Ann. Soc. Entomol. Fr. (N.S.)* 37, 105–127.
- Lake, J.A., 1994. Reconstructing evolutionary trees from DNA and protein sequences: paralogous distances. *Proc. Natl. Acad. Sci. USA* 91, 1455–1459.
- Lake, J.A., 1998. Optimally recovering rate variation information from genomes and sequences: pattern filtering. *Mol. Biol. Evol.* 15, 1224–1231.

- Larget, B., Simon, D., 1999. Markov chain Monte Carlo algorithms for the Bayesian analysis of phylogenetic trees. *Mol. Biol. Evol.* 16, 750–759.
- Lewis, P.O., 2001. A likelihood approach to estimating phylogeny from discrete morphological character data. *Syst. Biol.* 50, 913–925.
- Leache, A.D., Reeder, T.W., 2002. Molecular systematics of the Eastern Fence Lizard (*Sceloporus undulatus*): a comparison of parsimony, likelihood, and Bayesian approaches. *Syst. Biol.* 51, 44–68.
- Lockhart, P.J., Steel, M.A., Hendy, M.D., Penny, D., 1994. Recovering evolutionary trees under a more realistic model of sequence evolution. *Mol. Biol. Evol.* 11, 605–612.
- Mallatt, J., Sullivan, J., 1998. 28S and 18S rDNA sequences support the monophyly of lampreys and hagfishes. *Mol. Biol. Evol.* 15, 1706–1718.
- Mallatt, J., Sullivan, J., Winchell, C.J., 2001. The relationships of lampreys to hagfishes: a spectral analysis of ribosomal DNA sequences. In: Ahlberg, P. (Ed.), *Major Events in Early Vertebrate Evolution: Palaeontology, Phylogeny, and Development*. Taylor and Francis, London, pp. 106–118.
- Mallatt, J., Winchell, C.J., 2002. Testing the new animal phylogeny: first use of combined large-subunit and small-subunit rRNA gene sequences to classify the protostomes. *Mol. Biol. Evol.* 19, 289–301.
- Manuel, M., Kruse, M., Muller, W.E.G., Le Parco, Y., 2000. The comparison of beta-thymosin homologues among metazoa supports an arthropod–nematode link. *J. Mol. Evol.* 51, 378–381.
- Medina, M., Collins, A.G., Silberman, J.D., Sogin, M.L., 2001. Evaluating hypotheses of basal animal phylogeny using complete sequences of large and small subunit rRNA. *Proc. Natl. Acad. Sci. USA* 98, 9707–9712.
- Mittmann, B., 2002. Early neurogenesis in the horseshoe crab *Limulus polyphemus* and its implication for arthropod relationships. *Biol. Bull.* 203, 221–222.
- Murphy, W.J., Eizirik, E., O'Brien, S.J., Madsen, O., Scally, M., Douady, C.J., Teeling, E., Ryder, O.A., Stanhope, M.J., de Jong, W.W., Springer, M.S., 2001. Resolution of the early placental mammal radiation using Bayesian phylogenetics. *Science* 294, 2348–2351.
- Nardi, F., Spinsanti, G., Boore, J.L., Carapelli, A., Dallai, R., Frati, F., 2003. Hexapod origins: monophyletic or paraphyletic? *Science* 299, 1887–1889.
- Nielsen, C., 1995. *Animal Evolution, Interrelationships of the Living Phyla*. Oxford University Press, Oxford, England.
- Nielsen, C., 2001. *Animal Evolution, Interrelationships of the Living Phyla*, second ed. Oxford University Press, Oxford, England.
- Ohnishi, H., 2000. Unpublished submission to GenBank, Accession number AB052895.
- Omilian, A.R., Taylor, D.J., 2001. Rate acceleration and long-branch attraction in a conserved gene of cryptic daphniid (crustacea) species. *Mol. Biol. Evol.* 18, 2201–2212.
- Peterson, K.J., Eernisse, D.J., 2001. Animal phylogeny and the ancestry of bilaterians: inferences from morphology and 18S rDNA gene sequences. *Evol. Develop.* 3, 170–205.
- Rambaut, A., Grassly, N.C., 2001. Seq-Gen Sequence Generator: An application for the Monte-Carlo simulation of DNA sequence evolution along phylogenetic trees. Version 1.2.3.
- Ramsköld, L., Chen, J.-Y., 1998. Cambrian lobopodians: morphology and phylogeny. In: Edgecombe, G.D. (Ed.), *Arthropod Fossils and Phylogeny*. Columbia University Press, New York, pp. 107–150.
- Regier, J.C., Shultz, J.W., 1997. Molecular phylogeny of major arthropod groups indicates polyphyly of crustaceans and a new hypothesis for the origin of hexapods. *Mol. Biol. Evol.* 14, 902–913.
- Regier, J.C., Shultz, J.W., 2001. Elongation factor-2: a useful gene for arthropod phylogenetics. *Mol. Phy. Evol.* 20, 136–148.
- Richter, S., 2002. The Tetracnata concept: hexapod–crustacean relationships and the phylogeny of Crustacea. *Organ. Diver. Evolut.* 2, 217–237.
- Sanderson, M.J., Driskell, A.C., Ree, R.H., Eulenstein, O., Langley, S., 2003. Obtaining maximal concatenated phylogenetic data sets from large sequence databases. *Mol. Biol. Evol.* 20, 1036–1042.
- Schmidt-Rhaesa, A., 1998. Phylogenetic relationships of the nematomorpha—a discussion of current hypotheses. *Zool. Anz.* 236, 203–216.
- Schmidt-Rhaesa, A., 2002. Are the genera of Nematomorpha monophyletic taxa? *Zool. Scr.* 31, 185–200.
- Schmidt-Rhaesa, A., Bartolomaeus, T., Lemburg, C., Ehlers, U., Garey, J.R., 1998. The position of the arthropoda in the phylogenetic system. *J. Morphol.* 238, 263–285.
- Schnare, M.N., Damberger, S.H., Gray, M.W., Gutell, R., 1996. Comprehensive comparison of structural characteristics in eukaryotic cytoplasmic large subunit (23S-like) ribosomal RNA. *J. Mol. Biol.* 256, 701–719.
- Scholtz, G., 2002. The Articulata hypothesis – or what is a segment? *Organ. Diver. Evolut.* 2, 197–215.
- Schram, F.R., Jenner, R.A., 2001. The origin of hexapoda: a crustacean perspective. *Ann. Soc. Entomol. Fr. (N.S.)* 37, 243–264.
- Shear, W.A., 1997. The fossil record and evolution of the myriapoda. In: Fortey, R.A., Thomas, R.H. (Eds.), *Arthropod Relationships*. Chapman & Hall, London, pp. 211–219.
- Shimodaira, H., Hasegawa, M., 1999. Multiple comparisons of log-likelihoods with applications to phylogenetic inference. *Mol. Biol. Evol.* 16, 1114–1116.
- Shultz, J.W., Regier, J.C., 2000. Phylogenetic analysis of arthropods using two nuclear protein-encoding genes supports a crustacean + hexapod clade. *Proc. R. Soc. London B* 267, 1011–1019.
- Smith, S., Overbeek, W.R., Woese, C.R., Gilbert, W., Gillevet, P.M., 1994. The genetic data environment: an expandable GUI for multiple sequence analysis. *CABIOS* 10, 671–675.
- Snodgrass, R.E., 1938. Evolution of the Annelida, Onychophora and Arthropoda. *Smithson. Misc. Collect.* 97, 1–159.
- Spears, T., Abele, L.G., 1997. Crustacean phylogeny inferred from 18S rDNA. In: Fortey, R.A., Thomas, R.H. (Eds.), *Arthropod Relationships*. Chapman & Hall, London, pp. 169–187.
- Sullivan, J., Markert, J.A., Kilpatrick, C.W., 1997. Phylogeography and molecular systematics of the *Peromyscus aztecus* species group (Rodentia: Muridae) inferred using parsimony and likelihood. *Syst. Biol.* 46, 426–440.
- Sullivan, J., Swofford, D.L., 2001. Should we use model-based methods for phylogenetic inference when we know that assumptions about among-site rate variation and nucleotide substitution pattern are violated? *Syst. Biol.* 50, 723–729.
- Suzuki, Y., Glazko, G.V., Nei, M., 2002. Overcredibility of molecular phylogenies obtained by Bayesian phylogenetics. *Proc. Natl. Acad. Sci. USA* 99, 16138–16143.
- Swofford, D.L., 2001. PAUP*. Phylogenetic analysis using parsimony (*and other methods). Version 4.0 beta 10. Sinauer Associates, Sunderland, MA.
- Swofford, D.L., Olsen, G.J., Waddell, P.J., Hillis, D.M., 1996. Phylogenetic inference. In: Hillis, D.M., Moritz, C., Mable, B.K. (Eds.), *Molecular Systematics*, second ed. Sinauer, Sunderland, MA, pp. 407–514.
- Swofford, D.L., Waddell, P.J., Huelsenbeck, J.P., Foster, P.G., Lewis, P.O., Rogers, J.S., 2001. Bias in phylogenetic estimation and its relevance to the choice between parsimony and likelihood methods. *Syst. Biol.* 50, 525–539.
- Takezaki, N., Figueroa, F., Zaleska-Rutczynska, Z., Klein, J., 2003. Molecular phylogeny of early vertebrates: monophyly of the agnathans as revealed by sequences of 35 genes. *Mol. Biol. Evol.* 29, 287–292.
- Turbeville, J.M., Pfeifer, D.M., Field, K.G., Raff, R.A., 1991. The phylogenetic status of arthropods, as inferred from 18S rRNA sequences. *Mol. Biol. Evol.* 8, 669–686.

- Wägele, J.W., Misof, B., 2001. On quality of evidence in phylogeny reconstruction: a reply to Zrzavy's defence of the 'Ecdysozoa' hypothesis. *J. Zool. Syst. Evol. Res.* 39, 165–176.
- Walossek, D., Müller, K.J., 1998. Early arthropod phylogeny in light of the Cambrian "Orsten" fossils. In: Edgecombe, G.D. (Ed.), *Arthropod Fossils and Phylogeny*. Columbia University Press, New York, pp. 185–232.
- Waloszek, D., Dunlop, J.A., 2002. A larval sea spider (Arthropoda: Pycnogonida) from the Upper Cambrian 'Orsten' of Sweden, and the phylogenetic position of pycnogonids. *Palaeontology* 45, 421–446.
- Wang, S., Zhao, M., Li, T., 2003. Complete sequence of the 10.3 kb silkworm *Attacus ricini* rDNA repeat, determination of the transcriptional initiation site and functional analysis of the intergenic spacer. *DNA Seq.* 14, 95–101.
- Wheeler, W.C., 1997. Sampling, groundplans, total evidence and the systematics of arthropods. In: Fortey, R.A., Thomas, R.H. (Eds.), *Arthropod Relationships*. Chapman & Hall, London, pp. 87–96.
- Wheeler, W.C., Whiting, M., Wheeler, Q.D., Carpenter, J.M., 2001. The phylogeny of extant hexapod orders. *Cladistics* 17, 113–169.
- Wilcox, T.P., Zwickl, D.J., Heath, T.A., Hillis, D.M., 2002. Phylogenetic relationships of the dwarf boas and a comparison of Bayesian and bootstrap measures of phylogenetic support. *Mol. Phy. Evol.* 25, 361–371.
- Wills, M.A., Briggs, D.E.G., Fortey, R.A., 1997. Evolutionary correlates of arthropod tagmosis: scrambled legs. In: Fortey, R.A., Thomas, R.H. (Eds.), *Arthropod Relationships*. Chapman & Hall, London, pp. 57–66.
- Wills, M.A., Briggs, D.E.G., Fortey, R.A., Wilkinson, M., Sneath, P.H.A., 1998. An arthropod phylogeny based on fossil and recent taxa. In: Edgecombe, G.D. (Ed.), *Arthropod Fossils and Phylogeny*. Columbia University Press, New York, pp. 33–106.
- Wilson, K., Cahill, V., Ballment, E., Benzie, J., 2000. The complete sequence of the mitochondrial genome of the crustacean *Penaeus monodon*: are malacostracan crustaceans more closely related to insects than to branchiopods? *Mol. Biol. Evol.* 17, 863–874.
- Winchell, C.J., 2001. Combining large- and small-subunit ribosomal RNA genes for phylogenetic comparison: robust results for the deuterostome animals and the elasmobranch fishes. M.S. thesis. Washington State University, Pullman.
- Winchell, C.J., Sullivan, J., Cameron, C.B., Swalla, B.J., Mallatt, J., 2002. Evaluating hypotheses of deuterostome phylogeny and chordate evolution with new LSU and SSU ribosomal DNA data. *Mol. Biol. Evol.* 19, 762–776.
- Zrzavy, J., Stys, P., 1997. The basic body plan of arthropods: Insights from evolutionary morphology and developmental biology. *J. Evol. Biol.* 10, 353–367.